



Colorectal Cancer Fact Sheet

Cancer that starts in either the colon or the rectum is called colorectal cancer. Colorectal cancer is the third most common cancer in women (after breast and lung cancer) and men (after prostate and lung cancer). In 2014, more than 96,800 people in the United States are estimated to be diagnosed with colorectal cancer, with approximately 50,310 deaths annually.¹ In Arizona, in 2011, there were 1,283 men and 1,119 women diagnosed with invasive colorectal cancer and 917 colorectal cancer deaths.² Half of all colorectal cancer cases in 2011 were diagnosed in late stage. A review of race/ethnicity found White Non-Hispanics have the lowest late stage diagnosis (48%) and Asian & Pacific Islanders have the highest late stage diagnosis (62%).³

Additional Information & Support Resources

FIT at Fifty HealthCheck Program
www.fitatfiftyhealthcheck.org

Colon Cancer Alliance/Undy5000
www.ccalliance.org

National Cancer Institute
www.cancer.gov/cancertopics/types/colon-and-rectal

American Cancer Society
www.cancer.org/Cancer/ColonandRectumCancer

Colorectal Cancer Control Program
www.cdc.gov/cancer/crccp

Fight Colon Cancer
www.fightcolorectalcaner.org

Screen 4 Colon Cancer
www.screen4coloncancer.org

Lynch Syndrome International
www.lynchcancers.com


National Colorectal Cancer Roundtable
80% by 2018
www.nccrt.org/about/80-percent-by-2018

Colorectal Cancer Risk Factors¹


- ✦ Aging and being over 50 years old
- ✦ Smoking
- ✦ Family history
- ✦ Genetic alterations
- ✦ Poor diet
- ✦ Having colorectal polyps
- ✦ Having Ulcerative Colitis or Crohn's Disease

2011 Arizona Colorectal Cancer

Age Adjusted Invasive Cancer Incidence Rate
Sex by Race/Ethnicity



Percent of Colorectal Cancer Cases Diagnosed in Late Summary Stage*



WNH: White Non-Hispanic WH: White Hispanic Blk: Black AI: American Indian AS&PI: Asian & Pacific Islander
* Late Summary Stage = regional and distant Stage combined

Symptoms of Colorectal Cancer¹

- ✦ Diarrhea or constipation
- ✦ Feeling that your bowel does not empty completely
- ✦ Finding blood in your stools
- ✦ Finding your stools are narrower than usual
- ✦ Gas pains or cramps, or feeling full or bloated
- ✦ Losing weight with no known reason
- ✦ Feeling very tired all the time
- ✦ Nausea or vomiting

Early Detection¹

SCREENING FOR COLORECTAL CANCER SHOULD BEGIN AT AGE 50.

Colorectal cancer typically does not present with symptoms until the disease is late stage.¹ It is important to know your family history and understand your recommended age to be screened. If you are experiencing symptoms *at any age*, talk to your doctor about getting screened.

The screenings below detect polyps, cancer, or other abnormal areas.

Immunochemical fecal occult blood test (iFOBT):

Detects tiny amounts of blood in the stool, that may be from polyps or cancers.

Sigmoidoscopy: Rectum and the lower part of the colon are examined with a lighted tube.

Colonoscopy: Entire colon is examined using a long, lighted tube.

Double-contrast barium enema: X-rays are taken of the colon and rectum after being given an enema with a barium solution and air is pumped into the rectum.



This factsheet is the result of a coordinated effort by the Arizona Department of Health Services (ADHS) Cancer Prevention and Control Programs (Arizona Cancer Registry, HealthCheck Programs, and Arizona Cancer Control Program).

Arizona
Cancer Prevention and
Control
More Arizonans Will Live Longer

We acknowledge the Centers for Disease Control and Prevention (CDC) for its support under cooperative agreements 5U58DP003858-02 awarded to the Arizona Cancer Registry and 5U58DP002034-04 awarded to the Arizona Fit at Fifty Health Check Program. This fact sheet's contents are solely the responsibility of the authors and do not necessarily represent the official views of the CDC.