

**Research and Professional Briefs**

# How Much of Racial/Ethnic Disparities in Dietary Intakes, Exercise, and Weight Status Can Be Explained by Nutrition- and Health-Related Psychosocial Factors and Socioeconomic Status among US Adults?

YOUFA WANG, MD, PhD; XIAOLI CHEN, MD, PhD

**ABSTRACT**

Large disparities exist in obesity and other chronic diseases across racial/ethnic and socioeconomic status (SES) groups in the United States. This study examined how much of racial/ethnic differences in diet, exercise, and weight status could be explained by nutrition- and health-related psychosocial factors (NHRPF) and SES among US adults. Nationally representative data of 4,356 US adults from the 1994-1996 Continuing Survey of Food Intakes by Individuals and the Diet and Health Knowledge Survey were used. NHRPF were assessed using 24 questions and related index scores. Dietary intakes were assessed using two nonconsecutive 24-hour dietary recalls. The US Department of Agriculture 2005 Healthy Eating Index was applied to evaluate diet quality. Body mass index was calculated based on self-reported weight and height. SES was assessed using education and household income. Americans with higher SES had better NHRPF and Healthy Eating Index scores. There were some small racial/ethnic differences in NHRPF, including making food choices and awareness of nutrition-related health risks. Multivariable linear and logistic regression models revealed some racial/ethnic differences in diet, exercise, and body mass index, but few of these disparities was explained by NHRPF, whereas SES explained some. The odds ratio of body mass index  $\geq 25$  for non-Hispanic blacks compared with whites decreased by 38% after SES was adjusted for. For exercise, we found a

smaller change (9.5%) in the racial/ethnic differences when controlling for SES. In conclusion, NHRPF may explain very few, but SES may contribute some of the racial/ethnic disparities in diet, exercise, and weight status in the United States.

*J Am Diet Assoc. 2011;111:1904-1911.*

In the United States, large health disparities exist across racial/ethnic and socioeconomic status (SES) groups regarding obesity and many other chronic diseases (1). A good understanding of the determinants of health disparities is critical to help develop related interventions to eliminate health disparities, which is a national priority (2,3). Some previous studies have suggested significant racial/ethnic differences in eating behaviors that are independent of SES (4). African Americans may face a more negative chronic health condition trajectory compared with their white counterparts (5,6). Racial/ethnic disparity in obesity in the United States may be attributed to racial/ethnic disparity in SES (5), although our previous research suggests ethnic, sex, and age differences in the SES-obesity associations (7-9). Furthermore, only a small proportion of the racial/ethnic differences in body weight may be explained by SES (7-9).

We and others hypothesize that the between-group differences in nutrition- and health-related psychosocial factors (NHRPF), including nutrition knowledge and beliefs (NKB), are important contributors to the large racial/ethnic and SES differences observed in US adults' dietary intakes, exercise, and obesity. Some previous research has found positive associations of good nutrition knowledge, self-efficacy, and beliefs with desirable diet and health outcomes (10-13). However, to our knowledge, no studies have examined whether NHRPF and SES could explain the ethnic disparities in diet, exercise, and obesity in the United States.

Using nationally representative data, we examined whether and how much of the racial/ethnic differences in US adults' dietary intakes, exercise, and weight status may be explained by NHRPF and SES. We also tested how much of the racial/ethnic disparities in weight status could be explained by the differences in diet and exercise participation.

*Y. Wang is an associate professor of international health and epidemiology and X. Chen is an assistant scientist, Department of International Health, Center for Human Nutrition, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD.*

*Address correspondence to: Youfa Wang, MD, PhD, Center for Human Nutrition, Room E2546, Johns Hopkins Bloomberg School of Public Health, 615 N Wolfe St, Baltimore, MD 21205. E-mail: ywang@jhsph.edu*

*Manuscript accepted: June 13, 2011.*

*Copyright © 2011 by the American Dietetic Association.*

*0002-8223/ \$36.00*

*doi: 10.1016/j.jada.2011.09.036*

**Table 1.** Nutrition and health-related psychosocial factors (NHRPF) and healthy eating index (HEI) score, by sociodemographic characteristics of US adults<sup>a</sup>

Characteristic	n	Nutrition Knowledge and Belief Score <sup>b</sup>		Nutrition Important <sup>c</sup>		Food Choice <sup>d</sup>		Awareness <sup>e</sup>		Better NHRPF <sup>f</sup>		Healthy Eating Index	
		Mean±SE <sup>g</sup>	P value	%±SE	P value	Mean±SE	P value	Mean±SE	P value	%±SE	P value	Mean±SE	P value
<b>Sex</b>													
Male	2,219	35.50±0.23	<0.001	52.83±1.73	<0.001	16.52±0.05	<0.001	5.72±0.05	<0.001	15.41±1.00	<0.001	50.69±0.36	<0.001
Female	2,137	37.70±0.17		69.25±1.36		16.97±0.04		6.11±0.04		26.82±1.34		53.85±0.5	
<b>Age (y)</b>													
20-34	1,165	35.56±0.26	<0.001	54.65±1.67	<0.001	16.52±0.07	<0.001	5.64±0.06	<0.001	13.96±1.22	<0.001	50.44±0.65	<0.001
35-49	1,507	37.04±0.20		62.09±1.59		16.86±0.05		6.04±0.04		23.42±1.63		52.16±0.47	
50-65	1,684	37.54±0.21		69.55±1.57		16.91±0.05		6.15±0.04		28.59±1.65		55.23±0.43	
<b>Race/ethnicity</b>													
Non-Hispanic white	3,285	36.67±0.19		58.40±1.20	<0.001	16.65±0.05	<0.01	6.04±0.04	<0.001	21.78±0.87		52.28±0.4	<0.001
Non-Hispanic black	512	36.34±0.32		71.39±3.26		17.26±0.13		5.58±0.10		20.96±3.06		47.60±0.55	
Hispanic	411	36.77±0.46		71.58±3.29		16.92±0.18		5.49±0.11		17.93±2.21		54.36±1.02	
Other	148	36.11±0.65		58.42±6.53		16.69±0.28		5.78±0.15		20.33±4.72		59.29±1.69	
<b>Education</b>													
<High school	731	35.71±0.37	<0.001	66.03±3.67		16.91±0.08		5.17±0.10	<0.001	10.72±1.18	<0.001	49.41±0.64	<0.001
High school	1,552	36.03±0.29		60.17±1.90		16.70±0.08		5.69±0.05		18.46±1.21		49.91±0.49	
>High school	2,073	37.20±0.17		60.78±1.17		16.75±0.06		6.24±0.03		25.43±1.21		54.50±0.41	
<b>Income</b>													
Low	1,079	36.18±0.29	<0.05	64.83±2.58		16.75±0.09		5.39±0.06	<0.001	12.90±1.49	<0.001	49.49±0.75	<0.001
Middle	1,543	36.40±0.23		61.94±1.45		16.75±0.06		5.85±0.05		20.07±1.60		50.97±0.46	
High	1,734	36.95±0.20		59.41±1.50		16.76±0.05		6.15±0.05		24.94±1.31		54.37±0.45	
<b>Comorbidity</b>													
0	2,830	36.28±0.22	<0.001	58.88±1.21	<0.001	16.70±0.04	<0.05	5.83±0.04	<0.001	18.94±1.14	<0.001	51.80±0.41	<0.001
1	951	37.27±0.18		64.63±1.88		16.86±0.07		6.12±0.05		24.89±1.95		52.94±0.59	
≥2	575	37.88±0.42		72.08±2.75		16.96±0.12		6.17±0.07		31.29±3.35		54.80±0.73	

<sup>a</sup>Controlled for complex survey design. Values are shown as mean (SE) or proportion (SE). P value from analysis of variance and/or  $\chi^2$  test for continuous and categorical variables, respectively.

<sup>b</sup>Instrument consisted of 11 questions scored as "To you personally, is it very important (score: 4), somewhat important (score: 3), not too important (score: 2), or not at all important (score: 1) to consume the following nutrients or foods at appropriate levels: salt/sodium, saturated fat, fiber, cholesterol, fruits and vegetables, sugar, dairy products, etc?" The higher the score, the better nutrition knowledge and beliefs (range 11-44).

<sup>c</sup>Subjects' consideration about the importance of nutrition at food purchase was assessed by asking "When you buy food, how important is nutrition?" Response "very important" was categorized as Nutrition Important group, whereas subjects with other responses served as the reference group.

<sup>d</sup>Food choice: Subjects took considerations of 6 key factors affecting food choices, including food safety, nutrition, price, refreshment, convenience, and taste. The higher the score, the more consideration in buying food (score range 6-24).

<sup>e</sup>Awareness: Subjects were aware of the number of 7 health problems caused by unhealthy eating behaviors, such as fat, cholesterol, sodium, fiber, and calcium. The higher the number, the more awareness of health risks.

<sup>f</sup>Summarized from high nutrition knowledge and beliefs (score  $\geq$  median), more food choice consideration (score  $\geq$  median), and awareness of health risks.

<sup>g</sup>SE=standard error.

## METHODS

### Study Design and Subjects

A nationally representative multistage stratified sample of 16,103 noninstitutionalized persons aged 0 to 90 years residing in the United States participated in the US Department of Agriculture (USDA) Continuing Survey of Food Intakes by Individuals (CSFII) 1994-1996 (14). Of the 16,103 CSFII 1994-1996 participants, 9,872 adults aged 20 years and older provided 1 to 2 days of dietary intake information in 24-hour recalls conducted 3 to 10 days apart. The 24-hour recalls utilized an automated five-stage multiple-pass approach (15). Demographic, socioeconomic, and lifestyle data were collected for all CSFII participants.

One adult aged 20 years or older per household who participated in CSFII completed the Diet and Health Knowledge Survey (DHKS). Each was asked about self-perceptions of the adequacy of intake levels of nutrients and other dietary components, awareness of diet-health relationships, perceived importance of following dietary guidance for specific nutrients and other dietary components, behaviors related to fat and food safety, knowledge about food sources of fats and cholesterol, and self-perceptions about weight status (16).

Of these, 5,765 completed the DHKS. Ninety DHKS participants who provided only one 24-hour recall were excluded. Also excluded were 1,319 participants older than age 65 years. These participants were excluded to help obtain a sample of relatively healthy individuals with no potentially special dietary needs. Our analyses included a final sample of 4,356 individuals (2,219 men and 2,137 women) aged 20 to 65 years.

### Key Measures and Study Variables

**NHRPF.** Study participants answered 24 questions regarding NHRPF. Based on their answers, we assigned scores for each question and then created related several index scores to summarized answers to a set of related questions for each category of NHRPF, respectively.

**Category 1: NKB.** Each subject was asked about his or her NKB using 11 questions. We calculated an overall NKB score (range 11 to 44) to summarize subjects' answers to these questions. A high total score indicated a better NKB.

**Category 2: Consideration of key factors affecting food choices.** Subjects were asked six questions to identify factors they considered important when buying foods, including safety, taste, ease of preparation (convenience),

**Table 2.** Comparisons of estimated racial/ethnic differences in diet quality, exercise participation, and weight status with and without adjustment for nutrition- and health-related psychosocial factors (NHRPF) or socioeconomic status (SES) of US adults

Characteristic	Racial/Ethnic Group <sup>a</sup>				R <sup>2</sup>
	Non-Hispanic Black		Hispanic		
<b>Comparison of with and without adjustment for NHRPF<sup>b</sup></b>					
<b>Linear regression models</b>	<b>β±standard error</b>	<b>% change<sup>c</sup></b>	<b>β±standard error</b>	<b>% change<sup>c</sup></b>	
<b>Healthy Eating Index</b>					
Model 1	-3.37±.80**	—	3.85±.92**	—	0.15
Model 1+Nutrition knowledge and beliefs <sup>d</sup>	-3.38±.77**	0.30	3.71±.87**	-3.64	0.16
Model 1+Nutrition important <sup>e</sup>	-3.82±.83**	13.35	3.26±.92**	-15.32	0.17
Model 1+Food choice <sup>f</sup>	-3.44±0.79**	2.08	3.77±.91**	-2.08	0.15
Model 1+Awareness <sup>g</sup>	-3.15±.75**	-6.53	4.00±.89**	3.90	0.16
Model 1+Intention to improve diet <sup>h</sup>	-3.48±.78**	3.26	3.56±.93**	-7.53	0.16
Model 1+Nutrition knowledge and beliefs+Nutrition important+Awareness	-3.59±.78**	6.53	3.35±.86**	-12.99	0.19
Model 1+NHRPF <sup>i</sup>	-3.47±.78**	2.97	3.76±.91**	-2.34	0.15
<b>Body mass index</b>					
Model 1	2.07±.41**	—	.71±.63	—	0.11
Model 1+Nutrition knowledge and beliefs <sup>d</sup>	2.07±.40**	0.00	.72±.63	1.41	0.11
Model 1+Nutrition important <sup>e</sup>	2.12±.40**	2.42	.77±.65	8.45	0.11
Model 1+Food choice <sup>f</sup>	2.08±.42**	0.48	.72±.63	1.41	0.11
Model 1+Awareness <sup>g</sup>	2.07±.40**	0.00	.71±.63	0.00	0.11
Model 1+Intention to improve diet <sup>h</sup>	2.13±.39**	2.90	.86±.62	21.13	0.12
Model 1+Nutrition knowledge and beliefs+Nutrition important+Awareness	2.11±.40**	1.93	.77±.65	8.45	0.11
Model 1+NHRPF <sup>i</sup>	2.08±.40**	0.48	.72±.63	1.41	0.11
<b>Logistic regression models</b>	<b>Odds ratio (95% confidence interval)</b>	<b>% change<sup>c</sup></b>	<b>Odds ratio (95% confidence interval)</b>	<b>% change<sup>c</sup></b>	
<b>High Healthy Eating Index (≥80th percentile)</b>					
Model 1	0.56 (0.31, 1.03)	—	1.74 (1.16, 2.62)*	—	
Model 1+Nutrition knowledge and beliefs <sup>d</sup>	0.55 (0.30, 0.99)*	-1.79	1.69 (1.14, 2.50)*	-2.87	
Model 1+Nutrition important <sup>e</sup>	0.51 (0.28, 0.96)**	-8.93	1.57 (1.03, 2.38)*	-9.77	
Model 1+Food choice <sup>f</sup>	0.55 (0.30, 1.01)	-1.79	1.70 (1.14, 2.52)*	-2.30	
Model 1+Awareness <sup>g</sup>	0.57 (0.31, 1.05)	1.79	1.78 (1.18, 2.68)*	2.30	
Model 1+Intention to improve diet <sup>h</sup>	0.55 (0.30, 0.99)*	-1.79	1.65 (1.09, 2.48)*	-5.17	
Model 1+Nutrition knowledge and beliefs+Nutrition important+Awareness	0.52 (0.28, 0.96)**	-7.14	1.57 (1.04, 2.37)*	-9.77	
Model 1+NHRPF <sup>i</sup>	0.55 (0.30, 1.00)*	-1.79	1.23 (1.15, 2.59)*	-29.31	
<b>Exercise participation</b>					
Model 1	0.69 (0.49, 0.97)*	—	0.84 (0.57, 1.24)	—	
Model 1+Nutrition knowledge and beliefs <sup>d</sup>	0.69 (0.49, 0.96)**	0.00	0.83 (0.55, 1.25)	-1.19	
Model 1+Nutrition important <sup>e</sup>	0.65 (0.46, 0.92)**	-5.80	0.81 (0.54, 1.21)	-3.57	
Model 1+Food choice <sup>f</sup>	0.67 (0.48, 0.93)**	-2.90	0.84 (0.57, 1.26)	0.00	
Model 1+Awareness <sup>g</sup>	0.70 (0.49, 0.98)*	1.45	0.84 (0.57, 1.25)	0.00	
Model 1+Intention to improve diet <sup>h</sup>	0.68 (0.48, 0.95)*	-1.45	0.83 (0.56, 1.22)	-1.19	
Model 1+Nutrition knowledge and beliefs+Nutrition important+Awareness	0.66 (0.46, 0.93)**	-4.35	0.81 (0.53, 1.23)	-3.57	
Model 1+NHRPF <sup>i</sup>	0.69 (0.49, 0.96)*	0.00	0.83 (0.56, 1.24)	-1.19	
<b>Body mass index ≥25</b>					
Model 1	2.35 (1.62, 3.39)**	—	1.48 (0.96, 2.27)	—	
Model 1+Nutrition knowledge and beliefs <sup>d</sup>	2.35 (1.63, 3.40)**	0.00	1.48 (0.96, 2.29)	0.00	
Model 1+Nutrition important <sup>e</sup>	2.38 (1.64, 3.47)**	1.28	1.50 (0.96, 2.34)	1.35	
Model 1+Food choice <sup>f</sup>	2.30 (1.57, 3.36)**	-2.13	1.46 (0.94, 2.24)	-1.35	
Model 1+Awareness <sup>g</sup>	2.35 (1.62, 3.39)**	0.00	1.48 (0.96, 2.29)	0.00	
Model 1+Intention to improve diet <sup>h</sup>	2.43 (1.70, 3.49)**	3.40	1.59 (1.02, 2.46)*	7.43	
Model 1+Nutrition knowledge and beliefs+Nutrition important+Awareness	2.38 (1.64, 3.47)**	1.28	1.51 (0.96, 2.37)	2.03	
Model 1+NHRPF <sup>i</sup>	2.34 (1.62, 3.38)**	-0.43	1.47 (0.96, 2.26)	-0.68	
<b>Comparison of with and without adjustment for SES<sup>j</sup></b>					
<b>Linear regression models</b>	<b>β±standard error</b>	<b>% change<sup>k</sup></b>	<b>β±standard error</b>	<b>% change<sup>k</sup></b>	
<b>Healthy Eating Index</b>					
Model 2	-4.80±.81***	—	2.29±.86*	—	0.125
Model 2+Income	-3.97±.84***	-17.29	3.09±.88**	34.93	0.136
Model 2+Education	-3.77±.73***	-21.46	3.40±.85***	48.47	0.156
Model 2+Income+Education	-3.38±.77***	-29.58	3.71±.87***	62.01	0.159

(continued)

**Table 2.** Comparisons of estimated racial/ethnic differences in diet quality, exercise participation, and weight status with and without adjustment for nutrition- and health-related psychosocial factors (NHRPF) or socioeconomic status (SES) of US adults (continued)

Comparison of with and without adjustment for SES <sup>1</sup>					
Linear regression models	$\beta \pm$ standard error	% change <sup>k</sup>	$\beta \pm$ standard error	% change <sup>k</sup>	
<b>Body mass index</b>					
Model 2	2.26 ± .40***	—	.90 ± .58	—	0.106
Model 2+Income	2.18 ± .41***	-3.54	.82 ± .60	-8.89	0.107
Model 2+Education	2.08 ± .40***	-7.96	.73 ± .62	-18.89	0.111
Model 2+Income+Education	2.07 ± .40***	-8.41	.72 ± .63	-20.00	0.111
Logistic regression models	Odds ratio (95% confidence interval)	% change <sup>k</sup>	Odds ratio (95% confidence interval)	% change <sup>k</sup>	
<b>High Healthy Eating Index (≥80th percentile)</b>					
Model 2	0.47 (0.26, 0.84)**	—	1.44 (1.01, 2.06)*	—	—
Model 2+Income	0.53 (0.29, 0.96)**	12.77	1.65 (1.14, 2.39)*	14.58	—
Model 2+Education	0.50 (0.28, 0.91)**	6.38	1.57 (1.07, 2.31)*	9.03	—
Model 2+Income+Education	0.55 (0.30, 0.99)*	17.02	1.69 (1.14, 2.50)*	17.36	—
<b>Exercise participation</b>					
Model 2	0.63 (0.45, 0.89)*	—	0.73 (0.49, 1.09)	—	—
Model 2+Income	0.66 (0.47, 0.92)*	4.76	0.76 (0.52, 1.13)	4.11	—
Model 2+Education	0.68 (0.48, 0.96)*	7.94	0.82 (0.54, 1.25)	12.33	—
Model 2+Income+Education	0.69 (0.49, 0.96)*	9.52	0.83 (0.55, 1.25)	13.70	—
<b>Overweight/obesity (body mass index ≥25)</b>					
Model 2	2.38 (1.66, 3.40)**	—	1.52 (1.02, 2.24)*	—	—
Model 2+Income	2.43 (1.69, 3.50)**	2.10	1.55 (1.03, 2.35)*	1.97	—
Model 2+Education	2.26 (1.58, 3.25)**	-5.04	1.44 (0.95, 2.19)	-5.26	—
Model 2+Income+Education	1.48 (0.96, 2.29)**	-37.82	1.48 (0.96, 2.29)	-2.63	—

<sup>a</sup>Non-Hispanic white served as the reference group.  
<sup>b</sup>Model 1 adjusted for survey year, sex, age, education, income, region, urbanization, comorbidity, and self-rated health.  
<sup>c</sup>Change in  $\beta$  or odds ratio (%) from Model 1.  
<sup>d</sup>Instrument consisted of 11 questions scored as “To you personally, is it very important (score: 4), somewhat important (3), not too important (score: 2), or not at all important (score: 1) to consume the following nutrients/foods at appropriate levels: salt/sodium, saturated fat, fiber, cholesterol, fruits and vegetables, sugar, dairy products, etc?” The higher the score, the better the nutrition knowledge and beliefs (range 11-44).  
<sup>e</sup>Subjects’ consideration about the importance of nutrition at food purchase was assessed by asking, “When you buy food, how important is nutrition?” Response “very important” was categorized as Nutrition Important group, whereas subjects with other responses served as the reference group.  
<sup>f</sup>Subjects took considerations of 6 key factors affecting food choices, including food safety, nutrition, price, refreshment, convenience, and taste. The higher the score, the more consideration in buying food (score range 6-24).  
<sup>g</sup>Subjects were aware of the number of 7 health problems caused by unhealthy eating behaviors, such as fat, cholesterol, sodium, fiber, and calcium. The higher the number, the more awareness of health risks.  
<sup>h</sup>Intention to improve diet was assessed by asking subjects if they thought about their current diet habits and if they would be willing to make changes: “The things that I eat and drink are healthy, and there is no reason for me to make changes.” Subjects who answered “strongly disagree” or “somewhat disagree” were grouped as Intention to Improve Diet and those reporting “strongly agree” or “somewhat agree” served as the reference group (no intention to improve diet).  
<sup>i</sup>NHRPF summarized from high nutrition knowledge and beliefs (score ≥ median), more food choice consideration (score ≥ median), and awareness of health risks.  
<sup>j</sup>Model 2 adjusted for survey year, sex, age, region, urbanization, comorbidity, self-rated health, and nutrition knowledge and beliefs.  
<sup>k</sup>Change in  $\beta$  or odds ratio (%) from Model 2.  
\* $P < 0.05$ .  
\*\* $P < 0.01$ .  
\*\*\* $P < 0.001$ .

how well the food keeps (freshness), nutrition, and food price. We computed a total score for considerations of food choices (range 6 to 24). The higher the score was, the more considerations of high food quality that the subject took.

Furthermore, we evaluated subjects’ consideration about the importance of nutrition at food purchase using one question. Subjects who answered “very important” were categorized as the Nutrition Important group. They were compared with the others.

Category 3: Awareness of nutrition-related health risks. Subjects were asked to respond “yes” or “no” to a question about whether or not they had heard about any health problems caused by unhealthy eating, such as eating too much fat, salt, cholesterol or sugar, but not enough fiber or calcium. This was evaluated by counting the number of seven health problems caused by unhealthy eating of which subjects were aware (range 0 to

7). The higher the number, the better awareness of health risks subjects had. Subjects were further categorized in the Awareness group if they were aware of all seven of these health risks, vs the other subjects who were less aware.

Category 4: Overall NHRPF score. We created several indexes to summarize these NHRPF: Those who had higher NKB (score greater than or equal to the median NKB score), more food choice consideration (food choice score greater than or equal to the median), and awareness of the seven nutrition-related health risks (Awareness group) were categorized as the Good NHRPF group, whereas the others were treated as the Poor NHRPF group. Note that we only counted the importance of nutrition once when creating the NHRPF index.

Category 5: Intention to improve diet. Subjects were also asked to identify whether they thought about their current diet habits and would be willing to make changes:

"The things that I eat and drink are healthy, and there is no reason for me to make changes." Their intention to make changes was assessed by using a 4-point Likert scale ("strongly disagree," "somewhat disagree," "somewhat agree," and "strongly agree"). For our analyses, subjects who answered "strongly disagree" or "somewhat disagree" were grouped as Intention to Improve Diet and those reporting "strongly agree" or "somewhat agree" served as the reference group (no intention to improve diet).

The Cronbach's  $\alpha$  coefficients were .86 for NKB (.86 for non-Hispanic whites or for blacks, .85 for Hispanics, and .87 for others), and .74 for awareness of nutrition-related health risks (non-Hispanic whites .74, non-Hispanic blacks or Hispanics .73, others .79). Overall, the internal consistency reliability of NHRPF measures in this study was acceptable.

**Dietary Intakes and Exercise Participation.** Based on responses that uncovered types of foods consumed during the 2 survey days along with their portion sizes, nutrient intakes were calculated by the USDA. Averaged dietary intakes of foods and nutrients from the 24-hour recalls were used. To assess the overall quality of diet, we applied the USDA's new 2005 Health Eating Index (HEI), which has been described elsewhere (17,18). The HEI was revised to reflect the 2005 Dietary Guidelines for Americans and consists of 12 components. HEI scores ranged from 0 to 100 (best). High diet quality was defined as HEI  $\geq$ 80th percentile.

Exercise participation was evaluated by asking one question: "How often do you exercise vigorously enough to work up a sweat?" Those who answered "rarely or never" were categorized as No Exercise Participation, whereas subjects who reported regularly exercising were treated as the Exercise Participation group.

**Weight Status.** Body mass index (BMI) was calculated based on self-reported weight and height. Overweight was defined as  $25 \leq \text{BMI} < 30$ ; and obesity,  $\text{BMI} \geq 30$ .

**Sociodemographic Characteristics.** Based on self-reported information, subjects were grouped as non-Hispanic white, non-Hispanic black, Hispanic, and others. SES was assessed using education and household income. Education was based on "<high school" (<12 years), "high school," and "< high school"; and household income levels were based on poverty income ratio 0% to 130% (ie, poor, eligible for food stamps), 131% to 350% (ie, middle income), and  $\geq 350\%$  (ie, high income).

**Other Covariates.** Other covariates included comorbidity (ie, subjects reported whether they had chronic diseases diagnosed by their doctors, including diabetes, high blood pressure, heart disease, cancer, osteoporosis, high blood cholesterol, and stroke); self-rated health (this was assessed based on the following question: "In general, would you say that your health is 1) excellent, 2) very good, 3) good, 4) fair, or 5) poor" [subjects with answers of "fair" or "poor" were categorized as being in the "fair/poor health" group]); survey year; geographic region (ie, Northeast, Midwest, South, and West); and degree of urbanization of the geographical area (ie, metropolitan statistical area—central city, suburban, and rural).

## Statistical Analysis

Our independent variables were NHRPF and SES, whereas dependent variables included HEI (continuous variable), high diet quality (categorical variable: HEI  $\geq$ 80th vs HEI <80th percentile) exercise participation (categorical variable: yes vs no), BMI (continuous variable), overweight/obesity (categorical variable:  $\text{BMI} \geq 25$  vs  $\text{BMI} < 25$ ), and obesity ( $\text{BMI} \geq 30$ ), respectively. Diet and exercise participation became covariates when we studied the changes in the racial/ethnic differences in weight status.

First, we evaluated internal consistency reliability of NHRPF measures (eg, NKB, food choices score, and awareness of nutrition-related health risk score) by calculating Cronbach's  $\alpha$  coefficient (19). The coefficients were .86 for NKB (.86 for non-Hispanic whites or blacks, .85 for Hispanics, and .87 for others), .55 for food choices (non-Hispanic whites or blacks .52, Hispanics 0.63, and others .63), and .74 for awareness of nutrition-related health risks (non-Hispanic whites .74, non-Hispanic blacks or Hispanics .73, and others .79). Overall, the internal consistency reliability of NHRPF measures was acceptable. Second, using the analysis of variance and  $\chi^2$  tests, we tested the differences in the NHRPF index and the individual NHRPF factors and HEI by sociodemographic characteristics.

Third, multivariable linear and logistic regression models were fit to examine whether and how much of the racial/ethnic differences in diet, exercise, and weight status could be explained by NHRPF or SES. In particular, we assessed how the estimated racial/ethnic effects (ie,  $\beta$  and  $R^2$  in linear regression models and odds ratio [OR] in logistic regression models) would change whether NHRPF and SES were included in the models vs not. The base model included the following variables: survey year, sex, age, education, income, region, urbanization, comorbidity, and self-rated health. In further analysis, we added NHRPF to the models to examine the racial/ethnic differences in diet, exercise, and BMI that could be explained by NHRPF. We suspected that the differences would become smaller after NHRPF was included.

Furthermore, using multivariable linear regression models, we tested how much of the racial/ethnic disparities in BMI could be explained by diet and exercise. By using normal weight ( $\text{BMI} < 25$ ) as the reference, we fit multinomial logistic regression models for overweight ( $\text{BMI} 25$  to  $29.9$ ) and obesity ( $\text{BMI} \geq 30$ ). We included survey year, sex, age, education, income, region, urbanization, comorbidity, self-rated health, and NKB in our base models. We then added HEI, energy, and exercise to the model to examine the changes in  $\beta$  or OR.

In all of the multivariate analysis, survey year, age, sex, region, degree of urbanization, chronic disease, and self-rated health were considered as potential confounders. All analysis was conducted using survey-related commands in SAS (version 9.2, 2008, SAS Institute Inc, Cary, NC) to take complex sampling design into account to produce nationally representative estimates and correct estimates of standard errors. Statistical significance was set at  $P < 0.05$ .

**Table 3.** Changes in the estimated racial/ethnic difference in weight status of US adults, with and without adjustment for diet and exercise participation<sup>a</sup>

Characteristic	Non-Hispanic Black		Hispanic		R <sup>2</sup>
	$\beta \pm$ standard error	% change <sup>b</sup>	$\beta \pm$ standard error	% change <sup>b</sup>	
<b>Linear regression models</b>					
<b>Body mass index</b>					
Model 1	2.07 $\pm$ .40*	—	.72 $\pm$ .63	—	0.11
Model 1 + Healthy Eating Index	2.04 $\pm$ .40*	-1.45	.75 $\pm$ .63	4.17	0.11
Model 1 + Energy	2.06 $\pm$ .41*	-0.48	.71 $\pm$ .63	-1.39	0.11
Model 1 + Exercise	2.06 $\pm$ .40*	-0.48	.71 $\pm$ .64	-1.39	0.11
Model 1 + Healthy Eating Index + Exercise	2.04 $\pm$ .40*	-1.45	.74 $\pm$ .64	2.78	0.11
Model 1 + Energy + Exercise	2.05 $\pm$ .41*	-0.97	.70 $\pm$ .63	-2.78	0.11
<b>Multinomial regression models</b>					
<b>Overweight (body mass index 25-29.9)</b>					
Model 1	1.90 (1.31, 2.75)**	—	1.53 (1.04, 2.250*)	—	
Model 1 + Healthy Eating Index	1.89 (1.30, 2.73)**	-0.53	1.54 (1.05, 2.260*)	0.65	
Model 1 + Energy	1.89 (1.31, 2.74)**	-0.53	1.54 (1.04, 2.26)*	0.65	
Model 1 + Exercise	1.90 (1.31, 2.77)**	0.00	1.54 (1.05, 2.26)*	0.65	
Model 1 + Healthy Eating Index + Exercise	1.89 (1.31, 2.74)**	-0.53	1.54 (1.05, 2.27)*	0.65	
Model 1 + Energy + Exercise	1.90 (1.31, 2.75)**	0.00	1.54 (1.05, 2.27)*	0.65	
<b>Obesity (body mass index <math>\geq</math>30)</b>					
Model 1	3.29 (2.09, 5.18)***	—	1.39 (0.71, 2.75)	—	
Model 1 + Healthy Eating Index	3.24 (2.09, 5.04)***	-1.52	1.43 (0.73, 2.80)	2.88	
Model 1 + Energy	3.27 (2.08, 5.15)***	-0.61	1.39 (0.71, 2.73)	0.00	
Model 1 + Exercise	3.25 (2.08, 5.08)***	-1.22	1.38 (0.70, 2.73)	-0.72	
Model 1 + Healthy Eating Index + Exercise	3.20 (2.07, 4.96)***	-2.74	1.41 (0.72, 2.78)	1.44	
Model 1 + Energy + Exercise	3.23 (2.06, 5.04)***	-1.82	1.38 (0.70, 2.72)	-0.72	
<sup>a</sup> Non-Hispanic white served as the reference group. Model 1 adjusted for survey year, sex, age, education, income, region, urbanization, comorbidity, self-rated health, and nutrition knowledge and beliefs.					
<sup>b</sup> Change in $\beta$ or odds ratio (%) from Model 1.					
* $P < 0.05$ .					
** $P < 0.01$ .					
*** $P < 0.001$ .					

## RESULTS AND DISCUSSION

Table 1 shows that women, older Americans, those with higher SES (high education or high income), and people with comorbidities had better NHRPF and HEI (all  $P < 0.001$ ). There were some but small racial/ethnic differences in NHRPF. Non-Hispanic whites reported better awareness of nutrition-related health risks (score  $6.04 \pm 0.04$ ) than non-Hispanic blacks ( $5.58 \pm 0.10$ ), Hispanics ( $5.49 \pm 0.11$ ), and other racial/ethnic group members ( $5.78 \pm 0.15$ ) ( $P < 0.001$ ). Overall, these NHRPF factors showed desirable associations with healthy eating as indicated by HEI (data not shown).

### How Much of the Racial/Ethnic Differences in Diet, Exercise, and BMI Could Be explained by NHRPF or SES?

In general, compared to non-Hispanic whites, non-Hispanic blacks had higher BMI and lower HEI, and were less likely to participate in exercise, whereas Hispanics had better HEI (data not shown). The racial/ethnic differences in diet and BMI changed little after controlling

for NHRPF (Table 2) but changed some if SES was controlled for (Table 2). Once SES was controlled for, the black–white differences in HEI became smaller, whereas the white–Hispanic differences became greater. Our linear regression models indicated that the black–white difference in BMI became smaller ( $\beta$   $2.07 \pm .40$  vs  $2.26 \pm .40$ ; percent change in beta was small,  $-8\%$ ) if SES were controlled for (eg, the subjects had similar SES), and our logistic regression models showed similar patterns for overweight and obesity (Table 2). After controlling for education and income in the multivariable models, the percent change in beta of HEI ( $-3.38 \pm 0.77$  vs  $-4.80 \pm 0.81$ , respectively) was 30% for non-Hispanic blacks vs whites. The OR of overweight and obesity (BMI  $\geq 25$ ) for non-Hispanic blacks compared with non-Hispanic whites changed by 38% after education and income were adjusted for in the model (OR [95% confidence interval] 1.48 [0.96 to 2.29] vs 2.38 [1.66 to 3.40]). For exercise participation, we found a smaller change (10%) in the racial/ethnic differences (non-Hispanic blacks vs whites) when controlling for SES (OR [95% confidence interval] 0.69 [0.49 to 0.96] vs 0.63 [0.45 to 0.89]).

## How Much of the Racial/Ethnic Differences in Weight Status Was Explained by Diet and Exercise?

The estimated racial/ethnic differences (non-Hispanic blacks vs whites) in BMI and obesity changed little (percent changes in  $\beta$  or OR were  $<-3\%$ ) after diet and exercise were controlled for (Table 3). Similar results were found for Hispanics vs whites. It was likely due to the fact that some overweight and obese subjects might have changed their diet and exercise (what was observed in the survey) and due to potential measurement errors.

Our study based on US nationally representative data showed several important findings that could help enhance the understanding of the complex factors that affect disparities in diet, exercise, and obesity across ethnic and SES groups. Our results provided mixed evidence regarding our original hypothesis that NHRPF could help explain some of the ethnic differences in diet, exercise, and obesity. We found that the racial/ethnic differences in diet and BMI changed little after controlling for NHRPF in our models.

SES played a more important role than NHRPF to explain the ethnic differences in these outcomes. In our study, the percentage change in HEI was 30% for non-Hispanic blacks vs whites after controlling for SES. The risk of overweight and obesity for non-Hispanic blacks comparing with non-Hispanic whites changed by 38% after controlling for education and income; for example, the black–white difference was reduced by 38%. These findings differ from findings from a recent study. The Exploring Health Disparities in Integrated Communities study conducted in southwest Baltimore, MD, compared to national data of National Health Interview Survey and found no racial/ethnic disparities in obesity among poor, urban women sharing the same social context (20). The study included 449 black women and 322 white women living in the same social context with similar low income. In our study, the SES factors only explained  $<40\%$  of the black–white difference in obesity.

Ours and previous findings suggest that disparities in obesity in the United States are not predominantly due to individual psychosocial differences regarding these selected NHRPF factors, but may be more affected by their broader social environments. Poor quality retail food environments in disadvantaged areas, in conjunction with limited individual economic resources, contribute to increased risk of obesity within ethnic minorities and socioeconomically disadvantaged populations (21).

The  $R^2$  in our models shows that the many variables we included could only explain a very small proportion (10% to 20%) of the variations in BMI and HEI, which are likely due to the complex factors that affect BMI and HEI and due to potential measurement errors in these outcomes.

Two main limitations of this study are that only self-reported weight and height were collected in CSFII, and it is based on cross-sectional surveys. The study also has several important strengths, including that nationally representative data collected from a large sample with rich data regarding NHRPF were used; and we created a set of index scores to measure NHRPF and used the USDA new HEI to assess overall dietary quality.

## CONCLUSIONS

These US national survey data show that few of racial/ethnic differences in diet, exercise, and weight status were explained by health- and nutrition-related psychosocial factors, which is different from what we have expected, but SES explained considerable proportion; for example, 30% of the black–white difference in dietary quality and about 40% in overweight and obesity prevalence. However, the reduction in the white–black disparity for exercise participation was small (10%). The underlying causes of ethnic disparities in health-related behaviors such as eating and exercise and obesity in the United States are complicated. More well-designed studies with vigorous and comprehensive assessment of related factors are needed to help advance the understanding. Such knowledge is needed to help develop evidence-based national policies and programs to eliminate health disparities.

**STATEMENT OF POTENTIAL CONFLICT OF INTEREST:** No potential conflict of interest was reported by the authors.

**FUNDING/SUPPORT:** This study was supported by the National Institutes of Health/The National Institute of Diabetes and Digestive and Kidney Diseases (grant No. R01DK81335-01A1).

## References

1. Wang Y, Beydoun MA. The obesity epidemic in the United States—Gender, age, socioeconomic, racial/ethnic, and geographic characteristics: A systematic review and meta-regression analysis. *Epidemiol Rev.* 2007;29:6-28.
2. Agency for Healthcare Research and Quality. Key themes and highlights from the national healthcare disparities report. <http://www.ahrq.gov/qual/nhdr07/Key.htm>. Accessed May 24, 2010.
3. The Office of Disease Prevention and Health Promotion, US Department of Health and Human Services. Healthy People 2010. <http://www.healthypeople.gov/>. Accessed May 4, 2010.
4. Bahr PR. Race and nutrition: An investigation of black-white differences in health-related nutritional behaviours. *Sociol Health Illn.* 2007;29:831-856.
5. Kahng SK. Can racial disparity in health between black and white americans be attributed to racial disparities in body weight and socioeconomic status? *Health Soc Work.* 2010;35:257-266.
6. Farmer MM, Ferraro KF. Are racial disparities in health conditional on socioeconomic status? *Soc Sci Med.* 2005;60:191-204.
7. Zhang Q, Wang Y. Using concentration index to study changes in socio-economic inequality of overweight among US adolescents between 1971 and 2002. *Int J Epidemiol.* 2007;36:916-925.
8. Wang Y, Zhang Q. Are American children and adolescents of low socioeconomic status at increased risk of obesity? Changes in the association between overweight and family income between 1971 and 2002. *Am J Clin Nutr.* 2006;84:707-716.
9. Zhang Q, Wang Y. Trends in the association between obesity and socioeconomic status in US adults: 1971 to 2000. *Obes Res.* 2004;12:1622-1632.
10. Laforge RG, Greene GW, Prochaska JO. Psychosocial factors influencing low fruit and vegetable consumption. *J Behav Med.* 1994;17:361-374.
11. Patterson RE, Kristal AR, White E. Do beliefs, knowledge, and perceived norms about diet and cancer predict dietary change? *Am J Public Health.* 1996;86:1394-1400.
12. Trudeau E, Kristal AR, Li S, Patterson RE. Demographic and psychosocial predictors of fruit and vegetable intakes differ: Implications for dietary interventions. *J Am Diet Assoc.* 1998;98:1412-1417.
13. Satia-Abouta J, Patterson RE, Kristal AR, Teh C, Tu SP. Psychosocial predictors of diet and acculturation in Chinese American and Chinese Canadian women. *Ethn Health.* 2002;7:21-39.
14. Tippet KS, Yasmin SC. *Design and Operation: The Continuing Survey of Food Intakes by Individuals and the Diet and Health Knowledge*

- Survey, 1994-1996*. Washington, DC: US Department of Agriculture, Agricultural Research Service; 1998. Nationwide Food Surveys Report No. 96-1.
15. Automated multiple-pass method. US Department of Agriculture, Food Surveys Research Group Web site. <http://www.ars.usda.gov/Services/docs.htm?docid=7711>. Accessed February 7, 2011.
  16. DHKS questionnaire. US Department of Agriculture, Food Surveys Research Group Web site: <http://www.ars.usda.gov/SP2UserFiles/Place/12355000/pdf/dhks.pdf>. Accessed February 7, 2011.
  17. Healthy Eating Index. US Department of Agriculture, Center for Nutrition Policy and Promotion Web site. <http://www.cnpp.usda.gov/HealthyEatingIndex.htm>. Accessed April 3, 2010.
  18. Beydoun MA, Powell LM, Wang Y. Reduced away-from-home food expenditure and better nutrition knowledge and belief can improve quality of dietary intake among US adults. *Public Health Nutr*. 2009; 12:369-381.
  19. Cronbach LJ. Coefficient alpha and the internal structure of tests. *Psychometrika*. 1951;16:297-334.
  20. Bleich SN, Thorpe RJ Jr, Sharif-Harris H, Fesahazion R, Laveist TA. Social context explains race disparities in obesity among women. *J Epidemiol Community Health*. 2010;64:465-469.
  21. Ford PB, Dzewaltowski DA. Disparities in obesity prevalence due to variation in the retail food environment: Three testable hypotheses. *Nutr Rev*. 2008;66:216-228.

 American Dietetic Association

Evidence Analysis Library®

For additional information on this topic, visit  
ADA's Evidence Analysis Library at  
[www.adaevidencelibrary.com](http://www.adaevidencelibrary.com)